



A Note on Singular Nonlinear Boundary Value Problems for the One-Dimensional p -Laplacian

HAISHEN LÜ AND CHENGKUI ZHONG

Department of Mathematics, Lanzhou University
Lanzhou, Gansu, 730000, P.R. China

(Received January 2000; accepted March 2000)

Communicated by R. P. Agarwal

Abstract—Existence theorems about the positive solution for the singular equation $(\varphi_p(y'))' + f(t, y) = 0$, $y(0) = y(1) = 0$ are established. The results are obtained by using a fixed-point theorem in cones. © 2000 Elsevier Science Ltd. All rights reserved.

Keywords—One-dimensional singular p -Laplace equation, Cone, Compact map, Fixed point.

1. INTRODUCTION

The boundary value problem for the one-dimensional p -Laplacian

$$\begin{aligned}(\varphi_p(y'))' + f(t, y) &= 0, & t \in (0, 1), \\ y(0) &= y(1) = 0,\end{aligned}\tag{1}$$

where $\varphi_p(s) = |s|^{p-2}s$, $p > 1$, has been studied extensively. For details, see, for example, [1–4]. The boundary value problem treated in the above-mentioned references is not able to possess singularity.

In [5], Wang and Gao considered the BVP

$$\begin{aligned}(\varphi_p(y'))' + k(t)f(y) &= 0, & t \in (0, 1), \\ y(0) &= y(1) = 0.\end{aligned}$$

In the article, they supposed that $f(u)$ is nonincreasing in $(0, +\infty)$. It was a critical condition in the proof. On the other hand, many authors have considered a particular case of (1) for $p = 2$, see [6–8],

$$\begin{aligned}y'' + f(t, y) &= 0, & t \in (0, 1), \\ y(0) &= y(1) = 0.\end{aligned}$$

They all assumed that f was monotone for y . Monotone condition is very important.

The authors are supported by the National Science Foundation of China (19971037). We thank D. O'Regan for his help.

This paper discusses the singular boundary value problem (1), but we eliminate the monotone condition. We obtain an existence theorem by using a fixed-point theorem in cones which no one has used to study p -Laplacian.

The following well-known lemma is very crucial in our arguments, see [9, p. 240].

LEMMA 1.1. *Let X be a Banach space, $K \subseteq X$ a cone in X . For $0 < \delta < r$, define $K_r = \{y \in K : |y| < r\}$. Assume that $F : \overline{K_r} \rightarrow K$ is a compact map such that*

- (a) $Fy \neq \lambda y, \forall |y| = r$, and $\lambda > 1$;
- (b) $Fy \neq \lambda y, \forall |y| = \delta$, and $\lambda < 1$;
- (c) $\inf\{|Fy| : |y| = \delta\} > 0$.

Then F has a fixed point in $\overline{K_r} \setminus K_\delta$.

COROLLARY 1.1. *$F, \overline{K_r}, K$ are defined as Lemma 1.1, if*

- (a) $|Fy| \leq |y| = r, \forall |y| = r$;
- (b) $|Fy| \geq |y| = \delta, \forall |y| = \delta$.

Then F has a fixed point in $\overline{K_r} \setminus K_\delta$.

To be precise, we write $C^+[0, 1] = C([0, 1], [0, \infty))$. $B(\theta, \alpha) = \{y \in C^+[0, 1] : \|y\|_0 < \alpha\}$. $B[0, 1] = \{y \in C^+[0, 1] : y(0) = y(1) = 0, y \text{ is convex}\}$. It is easily verified that $B[0, 1] \subset C^+[0, 1]$ is a cone. $B_\alpha = B(\theta, \alpha) \cap B[0, 1]$. $\partial B_\alpha = \partial B(\theta, \alpha) \cap B[0, 1]$.

2. MAIN RESULTS AND PROOF

THEOREM 2.1. *Suppose the following.*

- (H1) $f \in C((0, 1) \times [0, \infty), [0, \infty))$, $f(t, y) \leq p(t)q(y)$, where $q \in C([0, \infty), (0, \infty))$, $0 < \int_0^{1/2} \varphi_p^{-1}(\int_s^{1/2} p(r) dr) ds + \int_{1/2}^1 \varphi_p^{-1}(\int_{1/2}^s p(r) dr) ds < \infty$.
- (H2) $\exists \alpha > 0$, such that
 - (i)

$$\max \left\{ \int_0^{1/2} \varphi_p^{-1} \left(\int_s^{1/2} p(r) \max_{y \in [g_\alpha(r), \alpha]} q(y) dr \right) ds, \right. \\ \left. \int_{1/2}^1 \varphi_p^{-1} \left(\int_{1/2}^s p(r) \max_{y \in [g_\alpha(r), \alpha]} q(y) dr \right) ds \right\} \leq \alpha;$$

- (ii) $f(t, 0) \leq f(t, u)$, $0 < t < 1$, $0 \leq u \leq \alpha$. $\exists t_0 \in (0, 1)$, such that $f(t_0, 0) > 0$, where

$$g_\alpha(t) = \begin{cases} \alpha t, & 0 \leq t \leq \frac{1}{2}, \\ \alpha(1-t), & \frac{1}{2} \leq t \leq 1. \end{cases}$$

- (H3) $\exists 0 < \beta < \alpha$, such that

$$\min \left\{ \int_0^{1/2} \varphi_p^{-1} \left(\int_s^{1/2} \min_{y \in [g_\beta(r), \beta]} f(r, y) dr \right) ds, \int_{1/2}^1 \varphi_p^{-1} \left(\int_{1/2}^s \min_{y \in [g_\beta(r), \beta]} f(r, y) dr \right) ds \right\} \geq \beta.$$

Then BVP (1) has at least one solution $y \in C^1[0, 1]$, $\varphi_p(y') \in C(0, 1)$, and $\beta \leq \|y\|_0 \leq \alpha$.

For $\forall y \in \overline{B(\theta, \alpha)}$, define

$$x(t) := \int_0^t \varphi_p^{-1} \left(\int_s^t f(r, y(r)) dr \right) ds - \int_t^1 \varphi_p^{-1} \left(\int_t^s f(r, y(r)) dr \right) ds, \quad 0 < t < 1.$$

Clearly, $x(t)$ is continuous, nondecreasing in $(0, 1)$ and $x(0+) < 0 < x(0-)$. Thus, $x(t)$ has zeros in $(0, 1)$. Let A be a zero of $x(t)$ in $(0, 1)$. Then,

$$\int_0^A \varphi_p^{-1} \left(\int_s^A f(r, y(r)) dr \right) ds = \int_A^1 \varphi_p^{-1} \left(\int_A^s f(r, y(r)) dr \right) ds. \quad (2)$$

Define the operator

$$T : \overline{B(\theta, \alpha)} \rightarrow B[0, 1] \text{ by} \\ (Ty)(t) := \begin{cases} \int_0^t \varphi_p^{-1} \left(\int_s^A f(r, y(r)) dr \right) ds, & 0 \leq t \leq A, \\ \int_t^1 \varphi_p^{-1} \left(\int_A^s f(r, y(r)) dr \right) ds, & A \leq t \leq 1, \end{cases} \quad (3)$$

where A is satisfied (2).

LEMMA 2.1. *The operator T is compact from $\overline{B(\theta, \alpha)}$ to $B[0, 1]$.*

PROOF. We first show $T\overline{B(\theta, \alpha)}$ is bounded. Put

$$q_\alpha = \sup\{q(y) : y \in [0, \alpha]\}, \\ u_\alpha(t) := \begin{cases} \int_0^t \varphi_p^{-1} \left(\int_s^{A^*} p(r) dr \right) ds \varphi_p^{-1}(q_\alpha), & 0 \leq t \leq A^*, \\ \int_t^1 \varphi_p^{-1} \left(\int_{A^*}^s p(r) dr \right) ds \varphi_p^{-1}(q_\alpha), & A^* \leq t \leq 1, \end{cases}$$

where A^* is a zero of the function

$$x(t) := \int_0^t \varphi_p^{-1} \left(\int_s^t q(r) dr \right) ds - \int_t^1 \varphi_p^{-1} \left(\int_t^s q(r) dr \right) ds, \quad 0 < t < 1,$$

and

$$\begin{aligned} u(t) &= (Ty)(t), & \forall y \in \overline{B(\theta, \alpha)}, & \quad 0 < t < 1, \\ u_\theta(t) &= (T\theta)(t), & & \quad 0 < t < 1. \end{aligned}$$

So

$$f(t, 0) \leq f(t, y) \leq q(t)q_\alpha, \quad (4)$$

$$(\varphi_p(u'_\alpha))' = -p(t)q_\alpha, \quad t \in (0, 1), \quad (5)$$

$$(\varphi_p(u'))' = -f(t, y), \quad t \in (0, 1), \quad (6)$$

$$(\varphi_p(u'_\theta))' = -f(t, 0), \quad t \in (0, 1). \quad (7)$$

We can obtain that $u_\theta(t) \leq u(t) \leq u_\alpha(t)$, $t \in (0, 1)$. Otherwise, there would exist $t_0 \in (0, 1)$ at which $u(t_0) > u_\alpha(t_0)$, and hence, there would exist an interval (a, b) such that $u(t) > u_\alpha(t)$ in (a, b) and $u(a) - u_\alpha(a) = u(b) - u_\alpha(b) = 0$. Let $m = u(B) - u_\alpha(B)$ be the positive maximum of $u(t) - u_\alpha(t)$ on $[a, b]$. Then $B \in (a, b)$ and $u'(B) = u'_\alpha(B)$. Integrating both sides of the equalities (5) and (6) over $[s, B]$, $a < s < B$, we get

$$\begin{aligned} u'_\alpha(s) &= \varphi_p^{-1} \left(\varphi_p(u'_\alpha(B)) + q_\alpha \int_s^B p(r) dr \right), \\ u'(s) &= \varphi_p^{-1} \left(\varphi_p(u'(B)) + \int_s^B f(r, y(r)) dr \right). \end{aligned}$$

Integrating both sides of the above equality from a to B , we obtain

$$\begin{aligned} u_\alpha(B) - u_\alpha(a) &= \int_a^B \varphi_p^{-1} \left(\varphi_p(u'_\alpha(B)) + q_\alpha \int_s^B p(r) dr \right) ds, \\ u(B) - u(a) &= \int_a^B \varphi_p^{-1} \left(\varphi_p(u'(B)) + \int_s^B f(r, y(r)) dr \right) ds. \end{aligned}$$

Consequently, we are led to a contraction $0 < m = u(B) - u_\alpha(B) \leq 0$ using inequality (4). It is similar that $u_\theta(t) \leq u(t)$, $t \in (0, 1)$. Hence,

$$\|Ty\|_0 \leq \|u_\alpha\|_0 = u_\alpha(A^*), \quad \forall y \in \overline{B(\theta, \alpha)}. \quad (8)$$

We next show the equicontinuity of $T\overline{B(\theta, \alpha)}$ on $[0, 1]$. For any $\varepsilon > 0$, from the continuity of $u_\alpha(t)$ on $[0, 1]$, it follows that there is a $\delta_1 \in (0, 1/4)$ such that

$$u_\alpha(2\delta_1), \quad u_\alpha(1 - 2\delta_1) < \varepsilon.$$

If $(Ty)(A) < \varepsilon$, then for any $t_1, t_2 \in [0, 1]$,

$$|(Ty)(t_1) - (Ty)(t_2)| \leq |(Ty)(A) - (Ty)(0)| < \varepsilon.$$

If $(Ty)(A) > \varepsilon$, then $A \in [2\delta_1, 1 - 2\delta_1]$, and hence, for $t \in [\delta_1, 1 - \delta_1]$,

$$|(Ty)'(t)| = \left| \varphi_p^{-1} \left(\int_t^A f(r, y(r)) dr \right) \right| \leq \varphi_p^{-1}(q_\alpha) \varphi_p^{-1} \left(\int_{\delta_1}^{1-\delta_1} p(r) dr \right) = L.$$

Put $\delta_2 = \varepsilon/L$, then for $t_1, t_2 \in [\delta_1, 1 - \delta_1]$, $|t_1 - t_2| < \delta_2$,

$$|(Ty)(t_1) - (Ty)(t_2)| \leq |(Ty)'(\xi)| |t_1 - t_2| < L\delta_2 = \varepsilon,$$

where ξ lies between t_1 and t_2 . Set $\delta = \min\{\delta_1, \delta_2\}$. Then for $t_1, t_2 \in [0, 1]$, $|t_1 - t_2| < \delta$,

$$|(Ty)(t_1) - (Ty)(t_2)| < \varepsilon, \quad (9)$$

which shows that $T\overline{B(\theta, \alpha)}$ is equicontinuous on $[0, 1]$.

We next claim that $T : \overline{B(\theta, \alpha)} \rightarrow B[0, 1]$ is continuous. Assume that $\{y_n\}_{n=0}^\infty \subset \overline{B(\theta, \alpha)}$ and $y_n(t)$ converges to $y_0(t)$ uniform on $[0, 1]$. Hence, $\{(Ty_n)(t)\}_{n=1}^\infty$ is uniformly bounded and equicontinuous on $[0, 1]$. The Arzela-Ascoli Theorem tells us that there exist uniformly convergent subsequences in $\{(Ty_n)(t)\}_{n=1}^\infty$. Let $\{(Ty_{n(m)})(t)\}_{m=1}^\infty$ be a subsequence which converges to $v(t)$ uniformly on $[0, 1]$ and $\{A_{n(m)}\}_{m=1}^\infty$ converges to \bar{A} . In addition,

$$u_\theta(t) \leq (Ty_n)(t) \leq u_\alpha(t).$$

Put

$$[a, b] = \{t \in [0, 1] : u_\alpha(t) \geq \max u_\theta(t) > 0\}.$$

Then $[a, b] \subset (0, 1)$ and $\{A_n\} \subset [a, b]$, where A_n is the maximum point of $(Ty_n)(t)$ in $(0, 1)$. Thus,

$$\begin{aligned} (Ty_n)(A_n) &= \int_0^{A_n} \varphi_p^{-1} \left(\int_s^{A_n} f(r, y_n(r)) dr \right) ds \\ &\leq \varphi_p^{-1}(q_\alpha) \int_0^b \varphi_p^{-1} \left(\int_s^b p(r) dr \right) ds, \\ (Ty_n)(A_n) &= \int_{A_n}^1 \varphi_p^{-1} \left(\int_{A_n}^s f(r, y_n(r)) dr \right) ds \\ &\leq \varphi_p^{-1}(q_\alpha) \int_a^1 \varphi_p^{-1} \left(\int_a^s p(r) dr \right) ds. \end{aligned}$$

Notice that

$$(Ty_n)(t) = \begin{cases} \int_0^t \varphi_p^{-1} \left(\int_s^{A_n} f(r, y_n(r)) dr \right) ds, & 0 \leq t \leq A_n, \\ \int_t^1 \varphi_p^{-1} \left(\int_{A_n}^s f(r, y_n(r)) dr \right) ds, & A_n \leq t \leq 1. \end{cases}$$

Inserting $y_{n(m)}$ and $A_{n(m)}$ into the above and then letting $m \rightarrow \infty$, we obtain

$$v(t) = \begin{cases} \int_0^t \varphi_p^{-1} \left(\int_s^{\bar{A}} f(r, y_0(r)) dr \right) ds, & 0 \leq t \leq \bar{A}, \\ \int_t^1 \varphi_p^{-1} \left(\int_{\bar{A}}^s f(r, y_0(r)) dr \right) ds, & \bar{A} \leq t \leq 1, \end{cases} \quad (10)$$

and

$$v(\bar{A}) = \int_0^{\bar{A}} \varphi_p^{-1} \left(\int_s^{\bar{A}} f(r, y_0(r)) dr \right) ds = \int_{\bar{A}}^1 \varphi_p^{-1} \left(\int_{\bar{A}}^s f(r, y_0(r)) dr \right) ds. \quad (11)$$

Here we have applied Lebesgue's dominated convergence theorem. From the definition of T , we know that $v(t) \equiv (Ty_0)(t)$ on $[0, 1]$. This shows that each subsequence of $\{(Ty_n)(t)\}_{n=1}^\infty$ uniformly converges to $(Ty_0)(t)$. Therefore, the sequence $\{(Ty_n)(t)\}_{n=1}^\infty$ itself uniformly converges to $(Ty_0)(t)$. This means that T is continuous at $y_0 \in \overline{B(\theta, \alpha)}$. Therefore, T is continuous on $\overline{B(\theta, \alpha)}$ since $y_0 \in \overline{B(\theta, \alpha)}$ is arbitrary. Thus, the Arzela-Ascoli Theorem implies that $T : \overline{B(\theta, \alpha)} \rightarrow B[0, 1]$ is compact map. ■

PROOF OF THEOREM 2.1. It is easily verified that $y(t)$ is a positive solution. $\Leftrightarrow y$ is a fixed point of the map T in $C^+[0, 1]$,

$$(Ty)(t) := \begin{cases} \int_0^t \varphi_p^{-1} \left(\int_s^A f(r, y(r)) dr \right) ds, & 0 \leq t \leq A, \\ \int_t^1 \varphi_p^{-1} \left(\int_A^s f(r, y(r)) dr \right) ds, & A \leq t \leq 1. \end{cases}$$

By Lemma 2.1, $T : \overline{B}_\alpha \rightarrow B[0, 1]$ is a compact map. We next claim that T satisfies Corollary 1.1.

(i) $\forall y \in \partial B_\alpha$, $\|y\|_0 = \alpha$. If y is convex, $g_\alpha(s) \leq y(s) \leq \alpha$, $s \in [0, 1]$. (H2) implies that

$$\begin{aligned} \|Ty\|_0 &= \int_0^A \varphi_p^{-1} \left(\int_s^A f(r, y(r)) dr \right) ds = \int_A^1 \varphi_p^{-1} \left(\int_A^s f(r, y(r)) dr \right) ds \\ &\leq \max \left\{ \int_0^{1/2} \varphi_p^{-1} \left(\int_s^{1/2} p(r) \max_{y \in [g_\alpha(r), \alpha]} q(y) dr \right) ds, \right. \\ &\quad \left. \times \int_{1/2}^1 \varphi_p^{-1} \left(\int_{1/2}^s p(r) \max_{y \in [g_\alpha(r), \alpha]} q(y) dr \right) ds \right\} \leq \alpha. \end{aligned} \quad (12)$$

(ii) $\forall y \in \partial B_\beta$, $\|y\|_0 = \beta$. If y is convex, $g_\beta(s) \leq y(s) \leq \beta$, $s \in [0, 1]$. (H3) implies that

$$\begin{aligned} \|Ty\|_0 &= \int_0^A \varphi_p^{-1} \left(\int_s^A f(r, y(r)) dr \right) ds = \int_A^1 \varphi_p^{-1} \left(\int_A^s f(r, y(r)) dr \right) ds \\ &\geq \min \left\{ \int_0^{1/2} \varphi_p^{-1} \left(\int_s^{1/2} \min_{y \in [g_\beta(r), \beta]} f(r, y) dr \right) ds, \right. \\ &\quad \left. \times \int_{1/2}^1 \varphi_p^{-1} \left(\int_{1/2}^s \min_{y \in [g_\beta(r), \beta]} f(r, y) dr \right) ds \right\} \geq \beta. \end{aligned} \quad (13)$$

So, BVP (1) has at least one solution $y \in C^1[0, 1]$, $\varphi_p(y') \in C(0, 1)$, and $\beta \leq \|y\|_0 \leq \alpha$. ■

REMARK 2.1. The proof is not completed by using the Schauder fixed-point theorem because $f(t, y)$ is not monotone about y in $(0, \infty)$.

REMARK 2.2. For the particular case of $p = 2$, the condition $0 < \int_0^{1/2} \varphi_p^{-1}(\int_s^{1/2} p(r) dr) ds + \int_{1/2}^1 \varphi_p^{-1}(\int_{1/2}^s p(r) dr) ds < \infty$ is equivalent to $t(1-t)p(t) \in L^1[0, 1]$.

COROLLARY 2.1. Suppose (H1) satisfies and $f(t, y)$ is nondecreasing about y in $(0, \infty)$. In addition, assume

(H4) $\exists \alpha > \beta > 0$, such that

- (i) $\max\{\int_0^{1/2} \varphi_p^{-1}(\int_s^{1/2} f(r, \alpha) dr) ds, \int_{1/2}^1 \varphi_p^{-1}(\int_{1/2}^s f(r, \alpha) dr) ds\} \leq \alpha$,
- (ii) $\min\{\int_0^{1/2} \varphi_p^{-1}(\int_s^{1/2} f(r, g_\beta(r)) dr) ds, \int_{1/2}^1 \varphi_p^{-1}(\int_{1/2}^s f(r, g_\beta(r)) dr) ds\} \geq \beta$.
- (iii) $\exists t_0 \in (0, 1)$, such that $f(t_0, 0) > 0$.

Then BVP (1) has at least one solution $y \in C^1[0, 1]$, $\varphi_p(y') \in C(0, 1)$, and $\beta \leq \|y\|_0 \leq \alpha$.

REMARK 2.3. For the boundary value problem

$$\begin{aligned} (\varphi_p(y'))' + f(t, y) &= 0, & t \in (0, 1), \\ y(0) &= 0, & y'(1) = c, \end{aligned} \tag{14}$$

we have a similar result. It is omitted.

REFERENCES

1. C. De Coster, Pairs of positive solutions for the one-dimensional p -Laplacian, *Nonlinear Anal.* **23**, 669–681 (1994).
2. D. O'Regan, Some general existence principles and results for $(\phi(y'))' = q(t)f(t, y, y')$, $0 < t < 1$, *SIAM J. Math. Anal.* **24**, 648–668 (1993).
3. M.A. del Pino, P. Drábek and R. Manásevich, The Fredholm Alternative at the first eigenvalue for the one dimensional p -Laplacian, *J. Diff. Equ.* **151**, 386–419 (1999).
4. R.F. Manasevich and F. Zanolin, Time-mappings and multiplicity of solutions for the one-dimensional p -Laplacian, *Nonl. Anal.* **21**, 269–291 (1993).
5. J. Wang and W. Gao, A singular boundary value problem for the one-dimensional p -Laplacian, *J. Math. Anal. Appl.* **201**, 851–866 (1996).
6. S. Taliaferro, A nonlinear singular boundary value problem, *Nonlinear Anal.* **3**, 897–904 (1979).
7. S. Taliaferro, On the positive solutions of $y'' + \phi(t)y^{-\lambda} = 0$, *Nonlinear Anal.* **2**, 437–446 (1978).
8. J. Janus and J. Myjak, A generalized Emden-Fowler equation with a negative exponent, *Nonlinear Anal.* **23**, 953–970 (1994).
9. K. Deimling, *Nonlinear Functional Analysis*, Springer-Verlag, (1985).
10. Q. Yao and H. Lü, Positive solutions of one-dimensional singular p -Laplace equation, (in Chinese), *Acta Mathematica Sinica* **41**, 1255–1264 (1998).